# DR. SCOTT LEVIN MD SOMERS ORTHOPEDIC SURGERY AND SPORTS MEDICINE GROUP

## Rotator Cuff Repair - Small/Medium

## **Phase 1 Protective Phase (0-6 Weeks)**

#### 0-6 Weeks

- Fit brace or sling (determined by physician).
- Begin pendulum exercises.
- Perform passive ROM to tolerance:

Flexion

ER/IR (shoulder at 45 degrees abduction)

- Perform elbow ROM.
- Perform hand-gripping exercises.
- Use ice and pain modalities.
- Discontinue brace or sling, either at 4 weeks or as per Dr. Levin
- Begin active assisted ROM exercises at 5 weeks

### Phase 2 Intermediate Phase (6-14 weeks)

#### Goals

- Establish full ROM (12 weeks)
- Gradually increase strength
- Decrease pain

#### 6-10 Weeks

- Perform active-assisted ROM and active ROM exercises as tolerated
- Begin isometric strengthening exercise
- Continue to work on grip and forearm strengthening
- Continue pain modalities.

## **10-14** weeks

• Begin isotonic and isokinetic strengthening exercises:

Deltoid to 90 degrees

ER/IR side-lying

Supraspinatus

Biceps/triceps

Scapular muscles

- Continue with active ROM exercises (full ROM by 12 to 14 weeks).
- Begin neuromuscular control exercises. If patient is unable to elevate arm with shoulder hiking (scapulothoracic substitution), maintain on humeral head stabilizing exercises.

## Phase 3-Advanced Strengthening Phase (14-26 Weeks)

#### Goals

- Maintain full, nonpainful ROM
- Improve strength of shoulder
- Improve neuromuscular control
- Gradual return to functional activities

#### 14-20 Weeks

- Continue active-assisted ROM exercise:
  - Flexion, ER, IR
- Perform self-capsular stretches.
- Begin aggressive strengthening program:

Shoulder flexion

Shoulder abduction (to 90 degrees)

Supraspinatus

ER/IR

Elbow flexors/extensors

Scapular strengthening

Begin conditioning program.

## **21-26 Weeks**

- Continue all exercises listed above.
- Begin interval sport program, if applicable.

## Phase 4 Return to Activity Phase (24 to 28 Weeks)

## Goals

• Gradual return to recreational sport activities

#### **24-28 Weeks**

- Continue all strengthening exercises.
- Continue all flexibility exercises.
- Continue progression on interval program.